

**REPORT DOCUMENTATION PAGE**Form Approved  
OMB NO. 0704-0188

Public Reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comment regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave Blank)		2. REPORT DATE 09/30/00	3. REPORT TYPE AND DATES COVERED <b>FINAL</b>
4. TITLE AND SUBTITLE Novel Coherent Effects in Semiconductors		5. FUNDING NUMBERS ARO Grant # DAAH04-96-1-0267	
6. AUTHOR(S) Nasser N. Peyghambarian-Principal Investigator; Rudolf H. Binder-Co-Principal Investigator		8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Arizona Optical Sciences Center 1630 East Univ. Blvd., Tucson, AZ 85721		10. SPONSORING / MONITORING AGENCY REPORT NUMBER <b>ARO 35642.1-PH</b>	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U. S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211			
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official department of the Army position, policy or decision, unless so designated by other documentation.			
12 a. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for public release; distribution unlimited.		12 b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) We have performed experimental and theoretical investigations of novel coherent effects in semiconductors, including excitonic Rabi oscillations, soliton-like pulse propagation at the exciton resonance, the nonlinear optical skin effect, interaction-induced polarization rotation in uniaxially strained quantum wells, light-induced ultrafast heavy-to-light hole population transfer, and the coherently coupled heavy-hole-light hole Stark effect. We have identified the role of semiconductor-specific aspects and contrasted the investigated effects with their two- and three-level counterparts. In general, we found semiconductor-specific aspects, such as the bandstructure and the Coulomb interaction, to modify these effects significantly.			
14. SUBJECT TERMS		15. NUMBER OF PAGES	
		16. PRICE CODE	
17. SECURITY CLASSIFICATION OR REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION ON THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT  UL

**20001122 096**

# **NOVEL COHERENT EFFECTS IN SEMICONDUCTORS**

**FINAL TECHNICAL REPORT**  
**August 1, 1995 through June 30, 2000**

**N. PEYGHAMBARIAN, PRINCIPAL INVESTIGATOR**  
**R. BINDER, CO-PRINCIPAL INVESTIGATOR**

**SUBMITTED:**  
**SEPTEMBER 30, 2000**

**U.S. ARMY RESEARCH OFFICE**

**(ARO Proposal Number: P-35642-PH)**  
**GRANT NUMBER: DAAHO4-96-1-0267**

**UNIVERSITY OF ARIZONA**  
**OPTICAL SCIENCES CENTER**  
**TUCSON, AZ 85721**

**APPROVED FOR PUBLIC RELEASE;**  
**DISTRIBUTION UNMLIMITED.**

**THE VIEWS, OPINIONS, AND/OR FINDINGS CONTAINED IN THIES REPORT ARE  
THOSE OF THE AUTHOR(S) AND SHOULD NOT BE CONSTRUED AS AN OFFICIAL  
DEPARTMENT OF THE ARMY POSITION, POLICY, OR DECISION, UNLESS SO  
DESIGNATED BY OTHER DOCUMENTATION.**

## **1. CO-PRINCIPAL INVESTIGATORS**

Nasser Peyghambarian (Principal Investigator)

Professor  
Optical Sciences Center  
University of Arizona

Rudolf H. Binder (Co-Principal Investigator)

Associate Professor  
Optical Sciences Center  
University of Arizona

## **2. STATEMENT OF THE PROBLEMS STUDIED**

During the time period covered by the ARO grant, we have investigated several coherent nonlinear effects in semiconductors. The investigations include both experimental and theoretical studies. They include various geometries (bulk semiconductors, semiconductor quantum wells, semiconductor micro cavities and vertical-cavity surface-emitting lasers) and a number of different optical excitation conditions. Common to all investigations is the issue of the ultrafast, coherent, nonlinear optical response of the semiconductor in the spectral vicinity of the fundamental band gap (especially the heavy-hole-exciton and light-hole-exciton response). Our studies included light-pulse propagation dynamics, vectorial polarization dynamics, and polarization-dependent excitonic population dynamics. The central issue common to all those individual studies was the role of semiconductor-specific many-body effects; in other words, the question of whether the observed and/or predicted effects can be understood in terms simple 2-level or 3-level dynamics. Quite generally, the answer to this question determines how future novel ultrafast optical semiconductor devices can be optimized in their performance.

Basically, we found in practically all areas significant deviations of the observed and/or predicted effects from simple 2-level or 3-level dynamics. The following is a list of individual aspects of this investigation and the main results obtained either experimentally, or theoretically, or through a combined theory-experiment analysis.

## **3. SUMMARY OF THE EXPERIMENTAL RESULTS**

1. We performed and reported an experimental study on coherent high intensity pulse propagation on the free exciton resonance in CdSe. Similar to the effect of self-induced transparency in atomic vapors our observations include pulse reshaping, pulse break-up, high transmission rates, and a minimum in the velocity dispersion for pulses tuned to the vicinity of the free exciton resonance. Self-induced transparency is a coherent pulse propagation phenomenon characterized by the almost unattenuated propagation of a pulse of  $2\pi$  area through several Beer's law absorption length. The observation of this effect requires the pulse duration to be smaller than the dephasing time of the relevant material excitation, otherwise incoherent pulse propagation occurs that is determined by linear absorption and, possibly, incoherent nonlinear absorption effects. Our system of choice are epitaxially grown CdSe crystals that exhibit an exciton resonance well separated from other bound and unbound electron-hole pair excitations with optical

thickness  $\propto L$  of 1.7 and 7, respectively. For the thinner sample the pulse transmission shows polariton beating as supported by model calculations. In the transmission of transform limited 180 fs pulses through the thicker sample we observe a temporal pulse break-up for intensities between 2 and 6 GW/cm<sup>2</sup> that provides a clear indication for coherent long distance propagation. At even higher intensities self-phase modulation causes characteristic temporal and spectral pulse break-up. Our studies show that coherent, low loss propagation of short optical pulses very close to an exciton absorption resonance in a semiconductor is indeed possible.

2. A theoretical study of soliton-like pulse propagation has been performed for both semiconductors and, as an additional investigation, for conventional nonlinear materials. In the case of semiconductors, we have found that certain previously known analytical solutions for solitons in conventional materials can also describe the soliton-like solution in semiconductors, although the underlying physics in both cases are completely different. Based on the one-dimensional propagation model of Maxwell's equation in the slow-varying envelope approximation, and the two-band semiconductor Bloch equations restricted to single-exciton inter-band polarizations, the temporal soliton-like character of moderate-intensity picosecond light pulses has been found to be dominated by excitonic exchange interactions. In contrast, phase-space blocking effects have been found to play only a minor role. Detailed comparisons with the phenomenon of self-induced transparency (SIT) have revealed that the important McCall-Hahn area theorem, well known for 2-level atoms, takes a completely different form in semiconductors. Whereas in 2-level systems the pulse area of the stable solutions is independent of the pulse duration, there is a distinct non-trivial relation between these two pulse characteristics in semiconductors.
3. We verified experimentally the dynamic nonlinear optical skin effect in a semiconductor. This effect leads to an internal reflection at a propagating boundary between areas of high and low exciton densities. To provide experimental proof of this theoretically predicted nonlinear optical effect, we studied in detail the properties of reflected light upon the incidence of intense femtosecond laser pulses onto a ZnSe crystal that is characterized by high optical nonlinearity at the fundamental exciton resonance. In a single beam experiment the dynamic nonlinear optical skin effect is manifested by an increasing amount of light that is reflected at photon energies extending to more than 20 meV below the energy of the exciton absorption resonance. The reflectivity far away from the exciton transition is unusually high and we obtained reflectivity values that are even larger than unity at photon energies four times the FWHM below the exciton resonance. A vivid explanation is that the frequency of the photons reflected at the (propagating) internal boundary between areas of high and low exciton densities is red-shifted due to the 'moving mirror' (Doppler) effect. In a second step, this frequency shift of the internally reflected light was used to study the propagation of the density front. Performing pump-probe reflection experiments with 150 fs pulses we showed that the Doppler-shifted reflection is present only for the duration of the pump pulse, indicating that the front propagation decelerates directly after the pump pulse leaves the sample. The experimental results agree very well with advanced model calculations that model the density front excitation and propagation deceleration within approximately 100 fs. We find that the density front excited by the strong fs laser pulse can propagate with a velocity of up to 10<sup>6</sup> m/s.
4. The following new coherent optical effect in semiconductors has been predicted: interaction -induced polarization rotation in uni-axially strained quantum wells. More precisely, we predict that high-intensity circularly-polarized ultra-short light pulses which are reflected at a uni-axially strained quantum well will induce a strong and time dependent change of ellipticity of the reflected light. This is caused by intrinsic exciton-exciton interactions. The theory on which these results are based is known as Hartree-Fock multi-band semiconductor Bloch theory.
5. In addition to basic ultrafast nonlinear optical effects in semiconductors, we have investigated vectorial polarization properties of VCSELs. First, we derived and classified the vectorial eigenmodes of VCSELs.

The core of the theory is a generalized vectorial transfer matrix method, and the basic ingredients are the cylindrical hybrid mode solutions of Maxwell's equations known for optical fibers. Combining the fiber mode theory with the generalized transfer matrix approach yields a complete characterization of optical mode frequencies, cavity-losses, longitudinal and transverse amplitude patterns of the electric and magnetic field in the VCSEL cavity, as well as the vector polarization characteristics of the fields. Furthermore, we have performed an investigation of the vectorial stability of linearly polarized fundamental modes in VCSELs with small amounts of anisotropic strain or stress in the optically active semiconductor quantum well. Our theory is a microscopic theory that includes the mutual dependence of dichroism and birefringence, and, based on this theory, we showed that for given nonzero strain one expects three pump current regimes, out of which two correspond to one stable linear polarization and one that corresponds to a bistable regime.

6. As a theoretical extension and application of the theory of dark states in semi-conductors, we have studied and predicted the light-induced adiabatic population transfer in semiconductor quantum wells. The population transfer under consideration involves heavy-hole and light-hole bands in p-doped semiconductor quantum wells. The investigation was based on an appropriate generalization of the multi-band semiconductor Bloch equations. In particular, the influence of many-body effects due to the Coulomb interaction between charge carriers in semiconductors has been studied. Based on our theory we predicted that this type of hole-population transfer is indeed possible, and we found that dynamic energy renormalizations due to Coulomb exchange interactions as well as quasi-thermalization of charge carriers significantly modifies the transfer dynamics as compared to that in simple 3-level systems.
7. Investigating the coherent response of a semiconductor quantum well to a strong resonant light field we reported the first experimental observation of multiple excitonic Rabi oscillations in a semiconductor material. Optical Rabi oscillations are amongst the most fundamental examples of coherent nonlinear light-matter interactions. In atomic and molecular two-level systems, optical Rabi oscillations are well established. Exposed to a strong stationary light field, the electron population oscillates between the lower and upper states with the Rabi frequency that is proportional to the dipole moment and the light field. Experimental difficulties arise from the fact that coherent processes in semiconductors are confined to much shorter times since the phases of the semiconductor excitations randomize over a characteristic time of hundreds of femtoseconds. Thus, experiments must be done with subpicosecond light pulses. Our experimental observation of excitonic Rabi oscillations in semiconductors was based on a unique pump-probe scheme. We excited the Rabi oscillations with a relatively long (770 fs) pump pulse and probed with a relatively short (150 fs) probe pulse to time gate the density oscillations. Utilizing pump and probe pulses that can be independently tuned in wavelength and temporally shaped we were able to create Rabi oscillations in the heavy-hole exciton density of an InGaAs quantum well and probe the transmission changes at the light-hole exciton transition. Since the light-hole exciton and the heavy-hole exciton both derive from the same conduction band oscillations in the heavy-hole exciton density manifest themselves in oscillations of the probe pulse absorption at the light-hole exciton resonance. We were also able to determine the dependence of the Rabi frequency upon pulse intensity. We found that the change in Rabi frequency with light intensity follows a square root behavior as in atomic two-level systems. However, modeling the experiment with the semiconductor Bloch equations shows that the Rabi frequency is renormalized due to Coulomb effects. Our results on this fundamental effect allowed us to analyze in a quantitative way analogies and principal differences in the coherent nonlinear response between two-level atoms and excitons in semiconductors.
8. Another extension of the theory of dark states and, more generally, 3-band dynamics is the area of 3-band Rabi oscillations. This is another novel coherent effect that we predicted theoretically. In contrast to 2-level and 2-band Rabi oscillations, 3-level and 3-band Rabi oscillations have been found to offer a rich variety of oscillation dynamics involving both, radiative and non-radiative (Raman) transitions. We have



analyzed excitonic 3-band Rabi oscillations in semiconductor quantum wells and found that the combination of band-coupling effects and Coulomb effects drastically modifies the 3-band Rabi oscillations as compared to those in simple 3-level systems.

9. We investigated experimentally and theoretically the coherently coupled optical Stark effect in a semiconductor three-state system under off-resonant excitation. This issue is a logical extension of the study of coherent exciton dynamics under resonant excitation condition (see above) that became possible by the granted six-month extension of the proposal. Semiconductor quantum wells that exhibit pronounced and well-separated heavy hole and light hole exciton resonances are in certain respects analogous to a three-state V-system. The optical response at the two resonances is coupled through a shared conduction band, as exhibited, for example, in our previous measurement of multiple excitonic optical Rabi oscillations. Here, we report on the observation of coherently coupled optical Stark shifts of heavy hole and light hole exciton resonances in an InGaAs multiple quantum well. The shifts are induced by 1.2 ps pulses pumping below the heavy hole exciton without excitation of a real exciton or carrier population. Since the splitting between heavy and light hole exciton resonance is even larger than the heavy hole exciton to pump photon energy detuning direct excitation of virtual light hole excitons is largely prohibit. This scheme enables us to study the heavy and light hole exciton dynamics solely induced by coherently driven virtual heavy hole excitons. We measured differential absorption spectra from weak, broadband probe pulses that reveal the dynamics of the semiconductor response at both heavy hole and light hole exciton resonance simultaneously. By employing both counter-circularly-polarized and co-circularly-polarized pump and probe, we were able to correlate shifts of heavy and light hole exciton states that share electron states with the same spin. In doing so, we have made the first measurement of the coherent coupling in a semiconductor three-band system under off-resonant excitation. The theoretical analysis shows that the leading contribution to the coherently coupled light hole exciton shift results from the shared electron state. Higher order Coulomb correlations play a minor rule as long as the pump energy is more than the biexciton binding energy below the heavy hole exciton resonance.

#### 4. PERSONNEL SUPPORTED

Dr. Rudolf H. Binder, Assistant Professor  
Dr. Axel Schülzgen, Assistant Research Professor  
Dr. Kai Wundke, Postdoctoral Research Associate  
Dr. Dariusz Burak, Postdoctoral Researcher  
Ilya Ruminantsev, Graduate Student  
Zhenghan Yang, Graduate Student  
Jason Auxier, PhD Student  
Oliver Hennig, MS, 1998  
Michael Donovan, PhD, scheduled for Oct. 2000  
George Mahs, PhD 1996

#### 5. PUBLICATIONS

- [1] "Shifted Self-Reflected Wave from a Semiconductor," A. Schülzgen, S. Hughes, and N. Peyghambarian, *Photonics West/Optoelectronics '97, San Jose (USA) 1997, Proceedings SPIE 2994*, 393 (1997).
- [2] "Observation of the Dynamic Nonlinear Optical Skin Effect in a Semiconductor," A. Schülzgen, S. Hughes, and N. Peyghambarian, *Invited Talk at CLEO/QELS '97, Baltimore (USA) 1997, OSA Technical Digest Series, Vol. 12*, 219 (1997).

- [3] "Green's function description of momentum orientation relaxation of photo-excited electron plasmas in semiconductors," R. Binder, H.S. Köhler, M. Bonitz, and N. Kwong, *Phys. Rev. B* **55**, 5110-5116 (1997).
- [4] "Cold-cavity vectorial eigenmodes of VCSELs," D. Burak and R. Binder, *IEEE J. Quant. Elect.* **33**, 1205-1214 (1997).
- [5] "Interaction-induced polarization rotation in anisotropic semiconductor quantum wells," R. Binder, , *Phys. Rev. Lett.* **78**, 4466-4469 (1997)
- [6] "Doppler-Shifted Internal Reflection of fs Pulses in a Semiconductor," A. Schülzgen and N. Peyghambarian, *Invited Talk at the 5th Int. Workshop on Femtosecond Technology, Tsukuba (Japan) 1998*.
- [7] "Doppler-Shifted Self-Reflection from a Semiconductor," A. Schülzgen, N. Peyghambarian, and S. Hughes, *phys. stat. sol. (b)* **206**, 125 (1998).
- [8] "Excitation Front Propagation in a Semiconductor induced by high Intensity Femtosecond Laser Pulses," A. Schülzgen and N. Peyghambarian, *Invited Talk at Photonics West/Optoelectronics '98, San Jose (USA) 1998, Proceedings SPIE 3283*, 740 (1998).
- [9] "Coherent High-Intensity Pulse Propagation on a Free Exciton Resonance in a Semiconductor," H. Giessen, S. Linden, J. Kuhl, A. Knorr, S. W. Koch, F. Grindele, M. Hetterich, M. Grün, S. Petillon, C. Klingshirn, and N. Peyghambarian, *Phys. Stat. Sol. (b)* **206**, 27 (1998).
- [10] "Semiconductor Kadanoff-Baym Equations Results for Optically Excited Electron-Hole Plasmas in Quantum Wells," N. H. Kwong, M. Bonitz, R. Binder, H.S. Köhler, *Phys. Stat. Sol. (b)* **206**, 197-201 (1998)
- [11] "Theoretical study of soliton-like propagation of picosecond light pulses interacting with Wannier excitons," I. Talanina, D. Burak, R. Binder, H. Giessen and N. Peyghambarian, *Phys. Rev. E* **58**, 1074-1080 (1998)
- [12] "Electromagnetic characterization of vertical-cavity surface-emitting lasers based on a vectorial eigenmode calculation," D. Burak and R. Binder, *Appl. Phys. Lett.* **72**, 891-893 (1998)
- [13] "A single-mode high-power vertical-cavity surface-emitting laser," T. Milster, W. Jiang, E. Walker, D. Burak, P. Claisse, P. Kelly, and R. Binder, *Appl. Phys. Lett.* **72**, 3425-3427 (1998)
- [14] "Identification of lasing modes of index-guided VCSELs under DC injection conditions," D. Burak, S.A. Kemme, R.K. Kostuk, and R. Binder, *Appl. Phys. Lett.* **73**, 3501-3503 (1998)
- [15] "Ultrafast adiabatic population transfer in p-doped semiconductor quantum wells," R. Binder and M. Lindberg, *Phys. Rev. Lett.* **81**, 1477-1480 (1998)
- [16] "Ultrafast adiabatic population transfer in p-doped semiconductor quantum wells," R. Binder and M. Lindberg, *contributed talk, in: Bulletin of the American Physical Society*, Vol. 43, No. 1, 765 (1998)

- [17] "Ultrafast adiabatic population transfer in p-doped semiconductor quantum wells," R. Binder and M. Lindberg, *contributed talk, paper QWE8, International Quantum Electronic Conference IQEC 1998, San Francisco, CA, May 3-8, 1998.*
- [18] "Theoretical Analysis of polarization properties of VCSEL eigenmodes," D. Burak and R. Binder, *contributed talk, Physics and Simulation of Optoelectronic Devices VI, OE '98, San Jose, California, 24-30 January 1998 (SPIE Proceedings Vol. 3283)*
- [19] "Full vectorial eigenmode analysis of VCSELs: threshold gain values and model frequencies," D. Burak and R. Binder, *contributed talk, Vertical Cavity Surface Emitting Lasers II, OE '98, San Jose, California, 24-30 January 1998 (SPIE Proceedings Vol. 3286)*
- [20] "Direct Observation of Excitonic Rabi Oscillations in Semiconductors," A. Schülzgen, R. Binder, M. E. Donovan, M. Lindberg, K. Wundke, H. M. Gibbs, G. Khitrova, and N. Peyghambarian, *Phys. Rev. Lett.* **82**, 2346 (1999).
- [21] "Experimental Observation of Multiple Excitonic Optical Rabi Oscillations in a Semiconductor," M. E. Donovan, A. Schülzgen, K. Wundke, R. Binder, M. Lindberg, H. M. Gibbs, G. Khitrova, and N. Peyghambarian, *Invited Talk at Photonics West/Optoelectronics '99, San Jose (USA) 1999, Proceedings SPIE 3624*, 216 (1999).
- [22] "Many-Body Aspects of Excitonic Rabi Oscillations in Semiconductors," R. Binder, M. Lindberg, A. Schülzgen, M. E. Donovan, K. Wundke, H. M. Gibbs, G. Khitrova, and N. Peyghambarian, *Invited Talk at Photonics West/Optoelectronics '99, San Jose (USA) 1999, Proceedings SPIE 3625*, (1999).
- [23] "Excitonic Rabi Oscillations in Semiconductor Quantum Wells," R. Binder, A. Schülzgen, M. E. Donovan, K. Wundke, H. M. Gibbs, G. Khitrova, N. Peyghambarian, and M. Lindberg, *Centennial Meeting of the American Physical Society, Atlanta (USA) 1999.*
- [24] "Excitonic Rabi Oscillations in Semiconductors," A. Schülzgen, R. Binder, M. E. Donovan, K. Wundke, H. M. Gibbs, G. Khitrova, N. Peyghambarian, and M. Lindberg, *Optics & Photonics News* **10**, 39 (1999).
- [25] "Laser-Induced Rabi Oscillations in Semiconductors," A. Schülzgen, R. Binder, M. E. Donovan, M. Lindberg, K. Wundke, H. M. Gibbs, G. Khitrova, and N. Peyghambarian, *Invited Talk at CLEO/QELS '99, Baltimore (USA) 1999, OSA Technical Digest Series*, 13 (1999).
- [26] "Rabi oscillations and Raman coherences in semiconductor quantum wells," R. Binder, *contributed paper QMC4, Quantum Electronics and Laser Science Conference (QELS '99), Baltimore Maryland, May 25-27, 1999*
- [27] "Microscopic model for polarization switching in optically anisotropic VCSELs," D. Burak, J. Moloney, and R. Binder, *Vertical Cavity Surface Emitting Lasers III, OE '99, San Jose, California, 25-26 January 1999 (SPIE Proceedings Vol. 3627)*
- [28] "Rabi Oscillations in Semiconductors," R. Binder, *29<sup>th</sup> Winter Colloquium on the Physics of Quantum Electronics (PQE), Snowbird, Utah, January 3-6, 1999*
- [22] "Experimental observation of multiple excitonic Rabi oscillations in a semiconductor," M. Donovan, A. Schülzgen, K. Wundke, R. Binder, M. Lindberg, H.M. Gibbs, G. Kithrova and N. Peyghambarian,



- [29] "Many-body aspects of excitonic Rabi oscillations in semiconductors," R. Binder, M. Lindberg, A. Schülzgen, M. Donovan, K. Wundke, H.M. Gibbs, G. Kithrova and N. Peyghambarian, *Physics and Simulations of Optoelectronic Devices VII, OE '99, San Jose, California, 25-29 January 1999 (SPIE Proceedings Vol 3625)*
- [30] "Laser-induced Rabi oscillations in semiconductors," A. Schülzgen, R. Binder, M. Donovan, M. Lindberg, K. Wundke, and N. Peyghambarian, contributed paper, *QMC5, Quantum Electronics and Laser Science Conference (QELS '99), Baltimore, Maryland, May 25-27, 1999.*
- [31] "A microscopical model for polarization stability in optically anisotropic VCSELs," D. Burak and R. Binder, *Conference on Physics and Simulation of Optoelectronic Devices, Photonics West 2000, San Jose, California, 22-28 January 2000 (SPIE Proceedings Vol. 3944)*
- [32] "Microscopic theory of polarization properties of optically anisotropic vertical-cavity surface emitting laser," D. Burak, J.V. Moloney, and R. Binder, *Phys. Rev. A* **61**, 53809-53830 (2000)
- [33] "A Green's function approach to the dynamics-controlled truncation formalism: I. Derivation of the  $\chi^{(3)}$  equations of motion," N. Kwong and R. Binder, *Phys. Rev. B* **61**, 8341-8358 (2000)
- [34] "Three-band excitonic Rabi oscillations in semiconductor quantum wells," R. Binder and M. Lindberg, *Phys. Rev. B* **61**, 2830-2836 (2000)
- [35] "Macroscopic vs. Microscopic description of polarization properties of optically anisotropic vertical-cavity surface-emitting lasers," D. Burak, J.V. Moloney and R. Binder, *IEEE J. Quant. Elect.* **36**, 956-970 (2000)
- [36] "On the identification of intervalence-band coherences in semiconductor quantum wells," R. Binder, I. Rummyantsev, N.H. Kwong and R. Takayama, *Phys. Stat. Sol. (b)* **221**, 169-174 (2000)
- [37] "Coherently Coupled Optical Stark Shifting in a Semiconductor Three-State System," M. E. Donovan, J. Lee, A. Schülzgen, I. Ruminantsev, R. Binder, and N. Peyghambarian, *to be submitted to Phys. Rev. B* (2000).